INDEX TO VOLUME XIII

New species in blackfaced type.

Abies, balsamea, Polystictus abietinus on, 184; grandis, 511; Polystictus abietinus on, 184; sp., host of Phomopsis pseudotsugae, 301

Abstracts, fourteenth annual meeting, American Phytopathological Society, 33–60; fourth annual meeting of Canadian Division, 290–294; first annual meeting of Inland Empire Section of Pacific Division, 509–512

Acer, rubrum, Polystictus hirsutus on, 185; saccharum, Polystictus pargamenus on, 185

Actinomyces, scabies, relation of soil moisture to, 231–236; chromogenus, in new land, 243

Aecidium nitens, 63, 69, 71

Agropyron, elongatum, host for zonate eye-spot, 59; intermedium, host for zonate eye-spot, 59; repens, host of bacterial disease, 76; repens, host for zonate eye-spot, 59; repens, host of Septoria agropyri, 7–8

Agrostis palustris, host of timothy rust, 38; stolonifera, host for zonate eye-spot, 59

Ailanthus, 185

Alfalfa, intracellular bodies in phloem tissue of, 326–329

Allard, H. A., Some possible relationships of the mosaic diseases, 555–557

Allium, sativum, host of Cercospora **Duddiae,** 362–365; cepa, host of Cercospora **Duddiae,** 362–365

Alnus tenuifolia, Polystictus hirsutus on, 185

Alternaria, 501–503; causing black point of wheat, 49; solani, 102; solani, reaction to fungicides, 532–542

American Phytopathological Society, fourth annual summer meeting, 150–152; fourteenth annual meeting, report, 188–198; Canadian Division, abstract of papers presented at the fourth annual meeting, 1922, 290–294; fifth annual summer meeting, 507; abstracts of papers presented at the Inland Empire Section, Pacific Division, Feb. 3, and 10, 1923, 509; Organization of the Inland Empire Section, Pacific Division, 511

Anchusa, capensis, host for aecial stage of leaf-rust of rye, 50; officinalis, host for aecial stage of leaf-rust of rye, 49

Anderson, P. J., Controlling tobacco wildfire in the seed-bed, 59; The relation of soil moisture to formaldehyde injury of onion seedlings, 392–403; and Osmun, A. V., An improved formaldehyde tank for the onion drill, 161–168

Anthostomella destruens, causing cranberry rot, 510

Anthracnose, of lettuce, notes on, 510

Aphids, importance of in overwintering mosaic, 555

Aphis, green peach, 511; green rose, 512; maidis, agent in transmission of sugarcane mosaic in Porto Rico, 24–29

Apothecia, photographing spore discharge, 30–32

Apple, bitter rot of, 290; bitter rot canker of, 461; blotch infection in 1922, 43; blotch, new aspects of control, 44; brown rot of, 462; canker on, due to Fusicoccum **pyrorum** n. sp., 225–230; cedar rust of, 303; fire blight of, 508; fruit rot due to Botrytis cinerea, 509; Phyllosticta leaf spot of, 290; Polystictus hirsutus on, 185; Polystictus lacteus on, 185; scab of, 508; scab control in Massachusetts, 44

Apricot, black heart disease due to Verticillium sp., 216–224; host of Coryneum beijerinckii, 510

Argentine Republic, curly leaf of sugar beets in, 458–460

Artichoke, globe, Botrytis rot, 58

Asparagus, dwarfing caused by Fusarium sp., 284; rust of, 300

Aspergillus, susceptibility of, to copper, 539

Avena, barbata, host of Leptosphaeria avenaria, 14; biologic specialization of Puccinia graminis avenae, 35; brevis, host of Leptosphaeria avenaria, 14;

fatua, host of Leptosphaeria avenaria, 14; nuda chinensis, host of Leptosphaeria avenaria, 14; sativa nigra, host of Leptosphaeria avenaria, 14; sativa orientalis, host of Leptosphaeria avenaria, 14; sterilis, host of Leptosphaeria avenaria, 14–15; strigosa, host of Leptosphaeria avenaria, 15

Bacillus, atrosepticus, relationship to Bacillus solanisaprus, 359–361; carotovorus, causing soft rot of Iris, 293; carotovorus, pectic fermentation by, 383–384; solanisaprus, relationship to Bacillus atrosepticus, 359–361

Bacteria, physiology of some plant pathogenic, 381–384

Bacterial organisms, two, causing leafspots on tobacco, 140–144

Bacterial spot of tomato and pepper, 307–315

Bacterium, angulatum, pectic fermentation, 383–384; campestre, pectic fermentation, 383–384; coronofaciens **atropurpureum** n. var., 81; exitiosum, on tomato and pepper, 308; malvacearum, new host of, 455–457; sojae, pectic fermentation, 383–384; solanacearum, on Chrysanthemum coronarium, 488–491; tabacum, pectic fermentation, 383–384; tumefaciens, 107–116; tumefaciens, on rose, 87–99; tumefaciens, on Paris daisy, 87–99; vesicatorium, on tomato and pepper, 308; on Solanum nigrum, 311; on Physalis minima, 311; on Datura stramonium var. tatula, 311; on Hyoscyamus niger and H. aureus, 311; on Nicotiana rustica, 311; on Solanum dulcamara, and S. rostratum, 311; on Lycium chinense and L. halimifolium, 311; on potato, 310; sp., pepper disease, 57

BAILEY, ALICE A., see LINK, GEO. L. L.

BAILEY, D. L., see STAKMAN, E. C.

BAILEY, IRVING W., Slime bodies of Robinia pseudo-acacia, 332–333

Banana wilt, 543; of Manila hemp plant, 253–256

Barberry, eradication by means of chemicals, 48; progress in eradication, 48; and Puccinia graminis in Denmark, 404–407

Barley, covered smut of, 153–160; inoculation with covered smut, 551–554; leaf rust of, 49; loose smut of, 153–160; new disease of in California, 562; Ophiobolus disease of, 562; rusty-blotch of in California, 562; Septoria leaf blotch of, 3–7; varietal susceptibility to covered smut, 552–553

BAXTER, D. V., White pine blister rust in Michigan, 285, 286

Bean, anthracnose of, 316–323; intracellular bodies in phloem tissue of, 326–329; sunscald, cause of, 376–380.

BEATTIE, R. KENT, see ORTON, W. A.

Begonia, host of Heterodera radicicola, 443

BENSAUDE, MATHILDE, A species of Olpidium parasitic in the roots of tomato, tobacco and cabbage, 451–454

Betula, occidentalis, Polystictus cinnabarinus on, 184; Polystictus hirsutus on, 185; lutea, Polystictus pargamenus on, 185

Bitter rot, 461

Bittersweet, host of Bacterium vesicatorium, 311

Blackberry, host of Kunkelia nitens, 508

Black heart disease, of apricots due to Verticillium sp., 216–224

Blackleg of potato, 359–361

Black point, morphology of durum wheat affected by, 48; some fungi causing, 49

Black rot, of grape, 462

Black stem rust, in Denmark, 404–407

BLAIR, R. J. and HALE, J. D. Wood v. fungi (Title only), 290

Blight, of canning peas, 293; of celery, 290; of cucurbits caused by Macrosporium cucumerinum, 199–204; of peony by Botrytis paeoniae, 292; peony, due to Botrytis cinerea, 509; tulip, 509; seedling blight of wheat and corn, 50

Blight, blossom, 509; of rose, 509; of snowball, 509; leaf, of golden seal, 509; of periwinkle, 509

Blister rust on Pinus cembra, P. monticola and P. strobus, 299
BLODGETT, F. M., Hot water and hot air treatment of potatoes, 55; Time-temperature curves for killing potato tubers by heat treatments, 465–475
Blossom blight, rose, 509; snowball, 509
Blotch, apple blotch infection in 1922, 43; of peony, by Cladosporium paeoniae, 292
Blue stem, of the black raspberry, 293
Bluestone, treatment of seed wheat and oats with, 169–183; treatment for smut, 512
BONCQUÉT, P. A., Discovery of curly leaf of sugar beets in the Argentine Republic, 458–460
Bordeaux mixture, potatoes sprayed with, 509; toxicity of different types, 532–542
BOTJER, J. CORTWIJN, and QUANJER, H. M., Dutch potato varieties resistant to wart disease and internal brown spot, 54
Botrytis, from apple, 462; blight of castor bean, 303; susceptibility of to copper, 539; allii, cause of onion neck rot, 509; cinerea, reaction to fungicides, 532–542; rot of globe artichoke, 58; in relation to damping-off of conifers, 387–391; reaction to staling products, 33–34; sulphur toxic to, 60; on apple, carrot, cranberry, fig, ginseng, gooseberry, lettuce, lilac, peony, pear, plum, raspberry, strawberry, squash, 509; paeoniae, on peony, 509; parasitica, on tulip, 509; sp., on cranberry, 510; on eggplant fruit, 509; on golden seal leaf, 509; on lettuce, 508; on periwinkle leaf, 509; on rose blossom, 509; on snowball blossom, 509
Bottom rot, of lettuce, 508
BOYLE, C., The growth reactions of certain fungi to their staling products, 33–34
Boyle, L. W., Fusarium rot of onions, 510; see HEALD, F. D.
Brachysporium, causing black point of wheat, 49
Breeding, nematode resistant plants, 436–450
BRENTZEL, W. E., A disease of flax not previously in the United States, 53–54; Disease of flax caused by a species of Rhizoctonia, 53
BRISLEY, HAROLD ROY, Studies on the blight of cucurbits caused by Macrosporium cucumerinum E. & E., 199–204
Broad bean, wilt of in fungous extract, 546, 547
Brome grass, Septoria leaf blotch of, 8–11
Bromus, altissimus, host of Septoria bromi, 9; carinatus, bacterial disease of, 78, 81; host of Bacterium coronafaciens var. atropurpureum, 75–86; inermis, bacterial disease of, 75; host of Bacterium coronafaciens var. atropurpureum, 75–86; host of Septoria bromi, 8–11, 17; host for zonate eyespot, 59; Septoria disease of, 8–11; secalinus, host of Septoria bromi, 9
BROWN, H. D., and GARDNER, MAX W., Lightning injury to tomatoes, 147
BROWN, J. G., and GIBSON, FREDERICK, A new host of Bacterium malvacearum, 455–457
BROWN, NELLIE A., Experiments with Paris daisy and rose to produce resistance to crown gall, 87–99
Bryophyllum, crown gall inoculations on, 476; studies on early stages of crown gall of, 476–482
Buffalo bur, host of Bacterium vesicatorium, 311
Bunt, control, 169–183; of wheat, treatment for, 511; occurrence in rye, 210–215
BURGER, O. F., Melanose and stem-end rot of citrus fruits, 45
Burgundy mixture, toxicity of, 532–542
BURKHOLDER, WALTER H., The gamma strain of Colletotrichum lindemuthianum (Sacc. & Magn.) B. et C., 316–323
Butternut, see Juglans cinerea

Cabbage, Chytridium brassicae on, 451; Olpidium brassicae on, 451–454; resistant to Fusarium, in 1922, 57; seed, hot water treatment for Phoma lingam, 251–253

Cacao, Meliola disease of, 247–250
Caeoma luminatum, 72
Calcium oxalate, crystals, in pustules of Melanconis juglandis, 418
Cancers, plant, 107–116
Cane blight, of raspberry, 508
Canker, due to Fusicoccum on apple, 225–230; Endothia parasitica, notes on occurrence of, 366–371; fig, due to Botrytis cinerea, 509; of Pseudotsuga, Larix, Tsuga, Abies, 301
Cantaloupe, host of Heterodera radicicola, 442; host of Macrosporium cucumerinum, 199–204; varieties susceptible to Heterodera radicicola, 444–446
Carica papaya, host of Heterodera radicicola, 442
Caron, O., Disinfection et parcelles (Title only, 290
Carrot, storage rot due to Botrytis cinerea, 509
Cartoons, in agricultural teaching, 33
Castanea dentata, 298; heart-rot of, 562; host of Polyporus spraguei, 288; Polystictus hirsutus on, 185; mollissima, 298; vesca, 298
Castor bean seedlings, crown gall inoculations on, 476; studies on early stage of crown gall of, 476–482
Catalpa, 185; speciosa, host of Heterodera radicicola, 442
Cedar, rust of, 303
Celery, blight of, 290; resistant to yellows, 56
Cephalothecium, in cotton seed, 501–503
Ceratostomella fimbriata, sweet potato black-rot fungus, 56
Cercospora, capsici, pepper disease, 57–58; **Duddiae** n. sp., causing leaf-spot on onion and garlic, 362–365; **lactucae** n. sp., causing leaf-spot of lettuce, 289
Cereals, leaf rust of, 338–340
Ceylon, Helminthosporium heveae in, 483
Chamaecyparis thyoides, Polystictus floridanus on, 184; Polystictus pinsitus on, 185; Poria papyracea on, 187
Chardon, C. E., and Veve, R. A., The transmission of sugar-cane mosaic by Aphis maidis under field conditions in Porto Rico, 24–29
Chaudhuri, H., see Paine, S. G.
Chemicals, eradicating the common barberry by means of, 48
Cherry, leaf-spot of, 508; host of Coryneum beijerinckii, 510; Polystictus hirsutus on, 185; Polystictus lacteus on, 185
Chestnut, bark disease, 298; resistance to bark disease, 47; treatment with lime sulphur, 366–371; with Bordeaux mixture, 366–371
Chlorophol, experiments with for control of barley smuts, 153–160
Christensen, J. J., see Dosdall, Louise
Chrysanthemum coronarium, host of Bacterium solanacearum, 488–491
Chupp, Charles, and Clapp, Grace L., Fusicoccum canker on apple, 225–230
Chytridium brassicae, on cabbage, 451
Cinnamomum camphora, relation of a Pestalozzia to disease of, 47–48
Citrus canker, 295
Citrus scab, history of in Japan, 492–495
Cladosporium citri, 495; elegans, 495
Clapp, Grace L., see Chupp, Charles
Clover, leaf-spot of, 59; powdery mildew of, 290
Coccomyces hiemalis, on cherry, 508
Colletotrichum, pepper diseases, 58; gossypii, sulphur toxic to, 60; lagenarium, 199; lindemuthianum, the gamma strain of, 316–323; trifolii, 121; sp., inoculations on cotton bolls, 502; isolated from interior of cotton seed, 501–502
Colley, Reginald H., and Taylor, Minnie W., Studies on the aecial stages of Cronartium ribicola and Cronartium occidentale, 46–47
Collins, J. Franklin, A noteworthy case of resistance to the chestnut bark disease, 47
Colloidal sulphur, 507
Conifers, damping-off of taproots of, 385–391
Coniothyrium diplodiella, on grapes, 518; fuckelii, 508

COOK, MEL. T., A disease of dahlias, 285; A greenhouse disease of melons, 462–463; Blister rust of the white pine, 283; Brown rot of apple, 462; Dwarf asparagus, 284; Early stages of crown gall, 476–482; The past and future of plant pathology (Title only), 290

COONS, G. H., Control of stinking smut of wheat (Tilletia levis) with dust treatment, 37; and NELSON, RAY, Yellows-Resistant celery: 3rd progress report, 56; see KOTILA, J. E.

Copper, acetate, toxicity of, 532–542; carbonate, dust for bunt control, 169–183; for oat smut control, 170; Corona compound of, 176; dust for smut control in wheat and oats, 293–294; experiments with for control of barley smuts, 153–160; results of tests with, 512; treatment for bunt, 511; sulphate, toxicity of, 532–542; and lime, for smut control, unsatisfactory, 294; treatment of seed wheat with, 169; anhydrous, 174, 177

Coriolus, abietinus, 294; balsameus, 294; delectans, 294; versicolor, 294

Corn, mildews, Sclerospora spontanea, S. maydis, S. sacchari, S. Philippinensis, 299; root, stalk and ear rot disease in Kansas, 52; relation of environment to infection by Diplodia zeae, 52–53; seedling blight of, 50

Corona compound of copper carbonate, 176

Corrosive sublimate, presprinkling before potato seed treatment with, 512

Corticium, vagum (Rhizoctonia solani), in relation to damping-off of conifers, 387–391; relation of temperature to the damping-off of garden cress seedlings by, 273–282

Coryneum, beijerinckii, on apricot, cherry, and peach, 510; blight of stone fruits in Washington, 510

Cotton, Pima-Egyptian, angular leaf-spot of, 455–457; seed, fungi isolated from the interior of, 501–503

COULSON, J. G., Peony diseases, 292–293; see DICKSON, B. T.

Coulure, of grapes, 513

Cowpea, Whippoorwill, nematode infection on, 439, 440

Cranberry, storage rots, 509–510; caused, by Anthostomella destruens, 510; by Botrytis sp., 510; by Fusicoccum putrefaciens, 510; by Guignardia vaccinii, 510; by Pestalozzia guepinii vaccinii, 510; by Phomopsis sp., 510; by Sporonema oxycocci, 510; by S. pulvinatum, 510; by Stemphylium sp., 510

CRAWFORD, R. F., Fungi isolated from the interior of cotton seed, 501–503

Cress, see Lepidium sativum

Crimson clover, see Trifolium incarnatum

Cronartium, occidentale, studies of aecia stages, 46–47; ribicola, foreign studies of, 45 infection through grafted roots, 46; in the Northwest, 46; on Pinus monticola 299; Pinus strobus, 299; Pinus cembra, 299; reaction to fungicides, 532–542; study of aecial stages, 46–47; wintering over test for, 283

CROWLEY, D. J., Preliminary report on the rots of cranberry in Pacific County, 509–510

Crown gall, early stages of, 476–482; location of organism in host tissues, 43; morphological responses of host tissues, 43; on tobacco, 107–116; method of inoculation, 107–108; inoculations, 109–115; see Bacterium tumefaciens

Cryptosphaerella viticola, on grapes, 518

Cryptothrips floridensis, causing thrips disease of camphor, 47

Cucumber, intracellular bodies in tissues of, 326–329

Cucumis melo, host of Heterodera radicicola, 442; sativus, host of Heterodera radicicola, 442

Cucurbita moschata, host of Heterodera radicicola, 442

Cucurbits, host of Macrosporium cucumerinum, 199–204

Culture, ascigerous stage of Fusarium moniliforme, 51

Cuprammonium sulphate, toxicity of, 532–542

Curly leaf, of sugar beets, 458–460

Currant, black, see Ribes nigrum; host for Fomes ribis, 508; tomato, host of Bacterium vesicatorium, 311
CUTLER, N. L., A contribution to the knowledge of the tree destroying fungi of the Vancouver forestry district, 294
Cynodon dactylon, host for zonate eye-spot, 59
Cytological studies of Tilletia tritici, 36–37
CZARNECKI, HELEN, Studies on the so-called black heart disease of the apricot, 216–224

Dactylis glomerata, host of timothy rust, 38
Daedalea confragosa, 294
Dahlias, disease of, caused by Sclerotinia libertiana, 285
Damping-off, relation to temperature in garden-cress seedlings, 273–282; of taproots of conifers, 385–391
DANA, B. F., Botrytis diseases in Washington, 509; Notes on Rhizoctonia, 509
Dandelion, host of Tylenchus dipsaci, 508
Datura, stramonium var. tabula, host of Bacterium vesicatorium, 311, 314
DAVIS, W. H., Germination of the spores of timothy smut (Ustilago striaeformis (Westd.) Niessl.), 38–39
Decay of living trees by Polystictus, 184–186
Degeneration, diseases of potato, 40
Denmark, severity of attacks of black stem rust in, 404–407
DETMERS, FREDA, Dothichiza canker on Norway poplar, 245–247
Diaporthe juglandis, ascogenous stage of Melanconium oblongum, 414, 420; disease of butternut, 47
DICKSON, B. T., Plant diseases of 1922 in western Quebec, 290; Temperature studies in mosaic diseases, 42; and HOOD, E. G., Temperature studies in mosaic diseases, 42; SUMMERBY, R., and COULSON, J. G., Control of oat smut, 291–292: see ECKERSON, S. H.
DICKSON, F., and FISHER, W. R., A method of photographic spore discharge from apothecia, 30–32
DICKSON, JAMES G., The influence of soil temperature and moisture on the development of seedling blight of wheat and corn caused by Gibberella saubinetii (Mont.) Sacc., 50; see JOHANN, HELEN
Dieback, gooseberry, due to Botrytis cinerea, 509
Diplodia, gossypii, 501–502; inoculations with, 502; zeae, Kansas corn disease, 52; relation of environment to infection by, 52–53
Disinfectants, organic mercury seed disinfectants, 38
Dissemination, of plant diseases, 295–306; in seeds, 303
DODGE, B. O., The distribution of the orange rust of Rubus, 61–74
DOOLITTLE, S. P., and MCKINNEY, H. H., Intercellular bodies in the phloem-tissue of certain plants and their bearing on the mosaic problem, 326–329
DORAN, W. L., Toxicity studies with some copper fungicides, 532–542
DOSDALL, LOUISE, and CHRISTENSEN, J. J., Variations in length of spores of Helminthosporium sativum P. K. & B. under different conditions of growth, 50
Dothichiza populea, on Populus eugenei, 245–247
Douglas fir, host of Fomes pinicola, 511
DOWELL, R. I., see STEVENS, F. L.
DRAYTON, F. L., Report of the Plant Disease Survey (Title only), 290
DRECHSLER, CHARLES, A new blossom-end decay of watermelons caused by an undescribed species of Pythium, 57; The occurrence of zonate eye-spot on various grasses and its mode of extension, 59
DUFF, G. H., Distribution of Ribes and Cronartium ribicola in Ontario (Title only), 293
DURAND, ELIAS JUDAH, Biography of, 408–409
DUSTAN, A. G., The natural control of the green apple bug (Lygus communis var. novascotiensis Knight) by a new species of Empusa, 290
Dusting, potatoes in 1922 in Mich., 54; of wheat and oats for smut, 169–183

Eastern blue stem of raspberry, 507, 508
ECKERSON, S. H., and DICKSON, JAMES G., The influence of soil temperature and moisture on the chemical composition of wheat and corn and their predisposition to seedling blight, 50–51; see MCKINNEY, H. H.
Egg plant, fruit rot due to Botrytis sp., 509
Elfingia megaloma, 294
ELLIOTT, JOHN A., The ascigerous stage of sweet potato black-rot fungus, 56; notice of death of, 464
Eleusine indica, host for zonate eye-spot, 59
Elymus virginicus, host for zonate eye-spot, 59
Empusa, the natural control of the green apple bug by a species of, 290
Endothia parasitica, 298; pregermination swelling of pycnospores of, 423; notes on occurrence and cankers of, 366–371
England, plant disease survey of, and Wales, 410
Eragrostis, major, host for zonate eye-spot, 59; pectinacea, host for zonate eye-spot, 59
Eutettix tenella, carrier of sugar beet curly leaf, 458–460
Eye-spot (zonate), on grasses, 59

Fagus atropunicea, Polystictus biformis on, 184
FAHMY, T., The production by Fusarium solani of a toxic excretory substance capable of causing wilting in plants, 543–550
False chinch bug, 512
Fermentation, pectic in culture media containing pectin, 381–384
FERNOW, KARL H., Spindling tuber or marginal leaf-roll, 40; A new host for potato mosaic, 40–41
Ficus carica, host of Heterodera radicicola, 442; canker, due to Botrytis cinerea, 509
Fig, see Ficus carica
Fir, Douglas, host of Fomes pinicola, 511; white, host of Fomes pinicola, 511
Fire blight, of apple, 508
FISHER, W. R., see DICKSON, F.
FITZPATRICK, H. H., A survey of the evidence indicating that Phytophthora should be merged with Pythium, 34
Flax, disease caused by Rhizoctonia, 53; a disease not previously reported in the United States, 53–54
FOLSOM, DONALD, see SCHULTZ, E. S.
Fomes, pinicola, causing wood decay, 511; ribis, on currant, 508; roseus, 294; ungulatus, 294; laricis, 294
Fomitiporella betulina, 294
Fomitiporia tsugina, 294
Foot-rot, of wheat, 561–562; varietal susceptibility of wheats to, 562
Formaldehyde, experiments with, for control of barley smuts, 153–160; as fungicide for wheat seed, 169; hot, presprinkling be ore potato seed treatment with, 512; improved tank for onion drill, 161–168; injury of onion seedlings, the relation of soil moisture to, 392–403; potato seed treatment with, 509; tank for onion drill, 58
Fourteenth annual meeting, American Phytopathological Society, 188–198
Fragaria chiloensis, host of Heterodera radicicola, 442
FRASER, W. P., and SIMMONDS, P. M., Dusting with copper carbonate and other substances for smut control, 293–294
Fraxinus americana, Polystictus pargamenus on, 185
Fruit rot, pear, plum, raspberry, strawberry, 509; of eggplant due to Botrytis sp., 509
Fungi, growth reactions to their staling products, 33–34; isolated from the interior of cotton seed, 501–503; relation of to mosaic virus, 557
Fungicide, sulphur as a, 60
Fungous filtrates, wilting power of, 549
Fusarium, cause of dwarf asparagus, 284; cultural reactions of, 291; influence of temperature on spores, 51; resistant cabbage in 1922, 57; rot of onions, 510; yellows of celery, 56; cubense, 543; in connection with wilt of Musa textilis, 253–256; moniliforme, ascigerous stage

in cultures, 51; pathogenicity of, on cereals, 52; presence in corn root, stalk and ear rot disease, 52; solani, toxic excretions of, 543–550; **spinaciae,** n. sp., Hungerford, 205–209; udum, cause of pigeon pea wilt, 543; vasinfectum, 463, 501; wilt of spinach, 205–209; spp., isolated from interior of cotton seed, 501–503; in relation to damping-off of conifers, 387–391

Fuscoporia ferruginosa, 294

Fusicladium effusum, on pecan, 300

Fusicoccum, canker on apple, 225–230; **pyrorum** n. sp., on apple, 225–230; putrefaciens causing cranberry rot, 510

GAINES, E. F., and STEVENSON, T. J., Occurrence of bunt in rye, 210–215

Galls, aerial galls of peach, 45

Ganoderma oregonense, 294; sessile, 294

Gardenia, nematode infected, 438, 442, 443

GARDNER, MAX W., and JACKSON, H. S., New aspects of apple blotch control, 44; and KENDRICK, JAMES B., Bacterial spot of tomato and pepper, 307–315; and KENDRICK, JAMES B., Field control of tomato mosaic, 372–375; see BROWN, H. D.

Garlic, leaf spot of, 362–365

Germination, Urocystis tritici, 134–137

Gibberella saubinetii, influence of temperature and moisture on blight of wheat and corn, 50; in Kansas, corn disease, 52

GIBSON, FREDERICK, see BROWN, J. G.

GILMAN, J. C., and MELHUS, I. E., Further studies on potato seed treatment, 341–358

Ginseng, moldy seed, due to Botrytis cinerea, 509; see Panax quinquefolium

Gloesporium, pepper diseases, 58; venetum, sulphur toxic to, 60

Gloeophyllum trabeum, 294; abietinellum, 294

Glomerella cingulata, 461; on grapes, 518; sulphur toxic to, 60; gossypii, 502; rufomaculans, 462

Gnomonia leptostyla, cause of leaf-spot of butternut, 411; perfect stage of Marsonia juglandis, 411

GODKIN, JAMES, see REDDY, CHAS. S.

Golden seal, leaf blight due to Botrytis sp., 509

Gooseberry, dieback due to Botrytis cinerea, 509

Grape, bitter-rot of, 518; black-rot of, 298, 300, 515; black-rot losses of, 516; blossom blight of, 515; dead-arm disease of, 518; downy mildew of, 298, 518; non-setting of, 513–519; pollen injury by rain, 515; phylloxera, 298; "rattles" disease of, 513; ripe-rot of, 518; self-sterility of, 514; shelling disease of, 513–519; white-rot of, 291, 518

GRAVES, ARTHUR HARMOUNT, The Melanconis disease of the butternut, 47; The Melanconis disease of the butternut (Juglans cinerea L.), 411–435

Green, peach aphis, 511; rose aphis, 512

Grey mold, lettuce, due to Botrytis cinerea, 509

GRIFFITHS, MARION A., see TISDALE, W. H.

Guignardia, bidwellii, blossom blight of grapes by, 513–519; vaccinii, causing cranberry rot, 510

Gum diseases of sugar cane in the Philippines, 504

Gymnoconia interstitialis, 61; reaction to fungicides, 532–542

Gymnosporangium clavipes, reaction to fungicides, 532–542

HALE, J. D., see BLAIR, R. J.

HANSEN, HANS R., Severity of attacks of black stem-rust in Denmark 1884 to 1921, 404–407

HARTER, L. L., LAURITZEN, J. I., and WEIMER, J. L., Mottle-necrosis of sweet potatoes, 145–146; Internal breakdown of sweet potatoes, 146–147; see LAURITZEN, J. I.

HARTGE, LENA, see HOTSON, J. W.

HARVEY, R. B., Biography of Elias Judah Durand, 408–409

HASKELL, R. J., Fourth summer meeting American Phytopathological Society,

150–152; Notice of international conference of phytopathologists in Holland, June 25 to 30, 1923, 287, 288

HEALD, F. D., Copper carbonate bunt problem, 511; and BOYLE, L. W., Further observations on the relation of the spore load to the per cent of bunt appearing in the crop (title only), 511; and BOYLE, L. W., Further notes on the relation of the spore load to the per cent of stinking smut appearing in the crop, 334–337; ZUNDEL, G. L., and BOYLE, L. W., The dusting of wheat and oats for smut, 169–183

Heart-rot, of chestnut, 562; of oaks, 562–563

Heat treatment, for potatoes, 465–475

Helianthus, annuus, taxonomy of Sclerotinia on, 496–499; debilis, host of Heterodera radicicola, 442; tuberosus, 499

Helminthosporium, **californicum** n. sp., on barley, 562; giganteum, zonate eyespot of grasses, 59; gramineum on barley, 562; **guianensis** n. sp., on Theabroma cacao, 249–250; heveae, in Ceylon, Java, Malay States, and South India, 483; in Sumatra, 483–487; on manihot, on papaya, and in rubber nurseries, 484; size of spores of, 485–487; oryzae, disease of rice, 53; sativum, causing black point, 49; on barley, 562; variations in length of spores, 50; sesameum, disease of Sesamum indicum, 53; teres, on barley, 562

Hemlock, western, host of Fomes pinicola, 511

HEMMI, TAKEWO, On the relation of temperature to the damping-off of garden cress seedlings by Pythium deBaryanum and Corticium vagum, 273–282

Henbane, see Hyoscyamus niger

HENRY, A. W., Some fungi causing black point of wheat, 49; The pathogenicity of Fusarium moniliforme Sheldon on cereals, 52

Heterodera radicicola, breeding plants resistant to, 436–450

Heterothallism, in Ophiobolus cariceti, 35

Hevea brasiliensis, host of Helminthosporium heveae, 483–487

HIGGINS, B. B., Seed treatment in the control of pepper diseases, 57–58

Hippeastrum, equestre, "mosaic" of, 42; johnsonii, intracellular bodies associated with a mosaic, 41–42

HOCKEY, J. F., Blue stem of the black raspberry, 293; The control of raspberry mosaic, 292

Hollyhock, rust on, 299

HOOD, E. G., see DICKSON, B. T.

HOPKINS, E. F., the Sphaerulina leaf-spot of clover, 59; 117–126

Hopper burn in Vermont, 237–241

Hordeum, deficiens, host of Septoria passerinii, 15; distichon, vars. erectum, nudum, nutans, hosts of Septoria passerinii, 15; horsfordianum, host of Septoria passerinii, 15; murinum, hybrid sterility of, 438; vulgare, Septoria leaf blotch of, 3–7; vulgare, vars. coerulescens, hexastrichum, himalaya, nigrum, pallidum, trifurcatum, hosts of Septoria passerinii, 15–16

Horse bean, see Vicia faba

HOTSON, J. W., and HARTGE, LENA, A disease of tomato caused by Phytophthora **mexicana**, sp. nov., 520–531

Hot water, experiments with, for control of barley smuts, 153–160; treatment of cabbage seed for Phoma lingem, 251–253

HOWARD, NATHANIEL O., The relation of an undescribed species of Pestalozzia to a disease of Cinnamomum camphora, 47

HOWITT, J. E., Two diseases new to Ontario, 291; and STONE, R. E., Abstract of five years' cooperative experiments in the prevention of oat smut, 292

HUNGERFORD, C. W., A Fusarium wilt of spinach, 205–209; A serious disease of wheat caused by Sclerotium rhizodes in Idaho, 463–464; Results of tests with copper carbonate in Idaho, 512; Preliminary results of experiments with leaf roll and mosaic in Idaho, 511, 512

HUNT, N. REX, Notes on the occurrence

and growth of cankers of Endothia parasitica, 366–371
Hyoscyamus, aureus, host of Bacterium vesicatorium, 311, 314; niger, host of Bacterium vesicatorium, 311, 314
Hypodermella laricis, causing leaf cast of Larix occidentalis in North Idaho, 505–506

Immunity of Prunus besseyi to aphis, 438
Inland Empire Section of the American Phytopathological Society, Pacific Division, organization of, 511
Inoculation, cross, with Heterodera radicicola, 442
Insects, as carriers of mosaic, 555–557; as agents for overwintering mosaic, 555–557
Internal break down of sweet potatoes, 146–147
International conference, Holland, 147–148
Intracellular bodies, in the phloem tissue of certain plants and their bearing on the mosaic problem, 326–329; associated with rosette of wheat, 41; associated with mosaic of Hippeastrum johnsonii, 41
Ipomoea, syringaefolia, host of Heterodera radicicola, 442; trichocarpa, Texas root rot, 33
Iris, bacterial soft rot of, 293

Jackson, H. S., see Mains, E. B.; see Gardner, Max W.
Java, Helminthosporium heveae in, 483
Johann, Helen; Influence of temperature on the morphology of Fusarium spores, 51; Dickson, James G., and Wineland, Grace, Relation of environments to infection of corn seedlings by Diplodia zeae (Schw.) Lev., 52
Johnson's mixture, toxicity of, 532–542
Johnston, C. O., Wheat smut investigations in Kansas: Report of progress 1920–21, 36; see Melchers, L. E.
Jones, Edith Seymour, Taxonomy of the Sclerotinia on Helianthus annuus, L. 496–499
Jones, L. R., Walker, J. C., and Tims, E. C., Work upon Fusarium-Resistant cabbage in 1922, 57
Juglans, cinerea, Melanconis disease of, 411–435; nigra, host of Melanconis juglandis, 413–415; regia, host of Melanconis juglandis, 415; host of Melanconium juglandinum in Europe, 414; sieboldiana, host of Melanconis juglandis, 415
Juniperus virginiana, Polystictus pinsitus on, 185

Kanred, susceptibility of F_2 hybrids with Marquis to stem rust, 437
Kansas, wheat smut investigations, 36
Kempton, F. E., Progress in barberry eradication, 48
Kendrick, James B., see Gardner, Max W.
Kentucky blue grass, Septoria leaf blotch, 11–13
Killough, D. T., see Taubenhaus, J. J.
Kirby, R. S., Heterothallism in Ophiobolus cariceti, 35
Kofoid, Charles A., Severin, Henry H. P., and Swezy, Olive, Nelson's spiral bodies in tomato mosaic not protozoans, 330–331
Kotila, J. E., and Coons, G. H., Spraying and dusting potatoes in 1922 at Chatham, Michigan, 54; Trypanosome-like bodies in solanaceous plants, 324–325
Krout, Webster S., Apple scab control in Massachusetts, 44
Kunkelia nitens, 61, 69; on blackberry, 508

Lactuca, sativa, host of Heterodera radicicola, 442; scariola, as host of Marssonia panattoniana, 510
Lactiporus speciosus, 294
Larch, leaf cast of, 505
Larix, occidentalis, leaf cast due to Hypodermella laricis, 505–506; sp., host of Phomopsis pseudotsugae, 301
La Rue, Carl D., Helminthosporium heveae in Sumatra, 483–487
Lasiagrostis splendens, host for zonate eyespot, 59

LAURITZEN, J. I., and HARTER, L. L., The relation of humidity to the infection of sweet potatoes by Rhizopus, 56; see HARTER, L. L.

Leaf blight, golden seal, periwinkle, due to Botrytis sp., 509; lilac, due to Botrytis cinerea, 509

Leaf cast, of Larix occidentalis, 505–506

Leaf curl, raspberry, 507–508

Leaf, curly, o sugar beets, 458–460

Leaf roll, potato, effect of heat treatment on, 465–475; potato, experiments with, in Idaho, 511; of potato in Northwest, 39, 40

Leaf rust of cereals, oversummering of, in Kentucky, 338–340

Leaf scorch, strawberry, 508

Leaf spot, angular, of tobacco, 140–144; of butternut, 411; of cherry, 508; of clover, 59, 117–126; disease of onion and garlic, 362–365; of peony, 290, 292; strawberry, 508; of apple, 290

Lee, H. ATHERTON, Gum diseases of sugar cane in the Philippines, 504; Sereh disease of sugar cane in Singapore, 145; and SERRANO, F. B., Banana wilt of the Manila hemp plant, 253–256

Leersia virginica, host for zonate eye-spot, 59

LEONIAN, LEON H., The physiology of perithecial and pycnidial formation in Valsa leucostoma, 257–272

Lepidium sativum, damping-off of, due to Pythium deBaryanum 273–282; due to Corticium vagum, 273–282

Leptosphaeria avenaria, on Avena barbata, A brevis, A fatua, A. nuda chinensis, A. sativa nigra, A. sativa orientalis, 14; A. sterilis, 14–15; A. strigosa, 15

Leptosphaeria coniothyrium, on raspberry, 508

Lettuce, anthracnose of, caused by Marssonia panattoniana, 510; grey mold due to Botrytis cinerea, 509; leaf spot of, due to Cercospora lactucae, 289; mosaic of, 104–106; mosaic, tip burn, bottom rot, drop, Botrytis rot, Botrytis sp., Rhizoctonia sp., Sclerotinia libertiana, 508; see Lactuca sativa

LEVINE, MICHAEL, Studies on plant cancers—V. Leafy crown galls on tobacco plants resulting from Bacterium tumefaciens inoculations, 107–116

LEVINE, M. N., and STAKMAN, E. C., Biologic specialization of Puccinia graminis secalis, 35; see STAKMAN, E. C.

Lightning injury to tomatoes, 147

Lilac, leaf blight due to Botrytis cinerea, 509

LINK, GEO. K. K., Mosaic and leaf roll of the potato in the Northwest, 39; RAMSEY, G. B., and BAILEY, ALICE A., Botrytis rot of the globe artichoke (Cynara scolymus), 58

Liquidambar styraciflua, Polystictus floridanus on, 184

Lotus corniculatus, host of Heterodera radicicola, 443

LUTMAN, B. F., An outbreak of hopper burn in Vermont, 237–241; Potato scab in new land, 241–244

Lycium, chinense, host of Bacterium vesicatorium, 311, 314; halimifolium, host of Bacterium vesicatorium, 311, 314

Lycopersicon esculentum, host of Heterodera radicicola, 442

Lycopersicum, esculentum, Phytophthora diseases of, 520; by P. mexicana, 520–531; pimpinnellifolium, host of Bacterium vesicatorium, 311

Lygus pratensis, 512

MACKIE, W. W., Foot-rot or Ophiobolus in California, 561–562; and PAXTON, G. E., A new disease of cultivated barley in California caused by Helminthosporium **californicum**, n. sp., 562

MACMILLAN, H. G., Cause of sunscald of beans, 376–380; Potato mosaic masking at high altitudes, 39

Macrosiphum solanifolii, 512

Macrosporium, pepper disease, 58; cucumerinum on cucurbits, 199–204; sarcinaeforme, sulphur toxic to, 60

MAINS, E. B., and JACKSON, H. S., Aecial stages of the leaf-rust of barley, Puccinia simplex, and of rye, P. dispersa, in the United States, 49; Strains of the leaf-rust of wheat, Puccinia triticina, in the United States, 36

Maize, attacked by Sclerospora maydis, 299; S. Philippinensis, 299; S. sacchari, 299; S. spontanea, 299; mildew of, 299

MAJOR, T. G., Cultural reactions of some dry root-rot organisms, 291

Malay States, Helminthosporium heveae in, 483

MALLOCH, W. S., The problem of breeding nematode resistant plants, 436–450

Mandarin, citrus scab on, 492

Manihot, Helminthosporium on, 484

Manila hemp plant, wilt of, 253–256

Marquis, susceptibility of F_2 hybrids with Kanred, 437

Marssonia panattoniana on lettuce, 510

Matrimony vine, host of Bacterium vesicatorium, 311

McCALLUM, A. W., The present status of white pine blister rust in Canada, 291

McCLINTOCK, J. A., Aerial galls of peach, 45

McCUBBIN, W. A., Plant pathology in public schools (Title only), 293

McINTOSH, G. E., Address by Dominion Fruit Commissioner (Title only), 290

McKINNEY, H. H., ECKERSON, SOPHIA H., and WEBB, R. W., Intracellular bodies associated with the rosette disease of wheat, 41; Intracellular bodies associated with a "mosaic" of Hippeastrum johnsonii, 41; see DOOLITTLE, S. P.

Medicago hispida reticulata, host of Heterodera radicicola, 443

Melanconis, carthusiana, relation to Melanconium juglandinum, 414; juglandis (E. and E.) comb. nov., description of, 415; disease of butternut, 47, 411; on Juglans nigra, 413; perfect stage of Melanconium oblongum, 412; sacchari, calcium oxalate crystals associated with, 418

Melanconium fuligineum, on grapes, 518; juglandinum, on Juglans regia, 414; relation to Melanconis carthusiana, 414; oblongum, acervuli of on butternut, 411; disease of butternut, 47; imperfect stage of Diaporthe juglandis, 414

Melanose of citrus fruits, 45

MELCHERS, L. E., and JOHNSTON, C. O., Corn root, stalk, and ear-rot disease investigations in Kansas: Report of progress 1922, 52

MELHUS, I. E., see GILMAN, J. C.

Meliola, disease of cacao, 247–250

Meliotus, alba, culture host for Ophiobolus cariceti, 35; **guianensis** n. sp., on cacao, 248–249

Melons, a greenhouse disease of, 462–463

Mercuric chloride, 470, 501; potato seed treatment with, 509

Mercury, organic, seed disinfectants, 38

METCALF, HAVEN, White pine blister rust in the Northwest, 46

Mildew, powdery, of clover, 290; raspberry, 508

Miscanthus japonicus, host of Sclerospora Philippinensis, 299

Moisture, influence on blight of wheat and corn, 50

Moldy seed, ginseng, due to Botrytis cinerea, 509

Mollisia earliana, 508

Mosaic, diseases, relationships of, 555–557; of sweet potato, 42; lettuce, 508; of lettuce, transmission by seeds, 104–106; occurrence of protozoa in, 41; new host for potato mosaic (Nicandra physaloides), 40; of peony, 292; potato mosaic, masking at high altitudes, 39; experiments with, in Idaho, 511; heat treatment for, 465–475; of potato in Northwest, 39; problem, intercellular bodies and their bearing on, 326–329; intracellular bodies associated with mosaic in Hippeastrum johnsonii, 41; raspberry, 507, 508; raspberry, control of by rogueing, 292; of sugar cane, 24–29; temperature studies, in mosaic diseases, 42; tomato, field control of, 372–375

Mottle-necrosis of sweet potatoes, 145–146

Mucor, racemosus, in relation to damping-off of conifers, 387–391
Muhlenbergia, mexicana, host for zonate eye-spot, 59; schreberi, host for zonate eye-spot, 59
Musa textilis, wilt of, 253–256
Muskmelon, see Cucumis melo
Mycosphaerella, fragariae, on strawberry, 508; rubina, Phoma stage on raspberries, 44
Myzus persicae on potato, 511

Neck rot, onion, 509
NELSON, RAY, The occurrence of protozoa in plants affected with mosaic and related diseases, 41; see COONS, G. H.
Nelson's spiral bodies in tomato mosaic not protozoans, 330–331
Nematode, breeding plants resistant to, 436–450; dandelion, 508
NEWHALL, ALLAN G., The importance of the Phoma stage of Mycosphaerella rubina causing spur blight of raspberries, 44; Seed transmission of lettuce mosaic, 104–106
Nicrandra phyaloides, a new host for potato mosaic, 40
Nicotiana, rustica, host of Bacterium vesicatorium, 311, 314; tabacum, host of Heterodera radicola, 442; var. Havanensis, tetramerous flowers in, 437; var. purpurea, lack of tetramerous flowers in, 437
Nicotine sulphate, 507
NOBLE, R. G., Studies on Urocystis tritici Koern., the organism causing flag smut of wheat, 127–139
Nonea rosea, host for aecial stage of leaf-rust of rye, 49
Norway poplar, Dothichiza populea on, 245–247
Nysius ericae, 512
Nyssa sylvatica, Polystictus pargamenus on, 185

Oats, dusting of, for smut, 169–183; smut, control of, by copper carbonate, 291–292
OCFEMIA, G. O., Helminthosporium disease of rice, 53
Oedocephalum, susceptibility of to copper, 539
Olpidium, brassicae, on cabbage, 451; sp. on roots of tomato, tobacco and cabbage, 451–454
Onion, formaldehyde tank for onion drill, 58; Fusarium, 508; Fusarium rot of, 510; leaf-spot of, 362–365; neck rot due to Botrytis allii, 509; pink root of, 290, 291, 508; seedlings, the relation of soil moisture to formaldehyde injury of, 392–403; smut, 508; treatment with formaldehyde, 161–168; Urocystis cepulae, 508
Onion drill, improved formaldehyde tank for, 161
Ophiobolus, cariceti, Heterothallism in, 35; graminis, on wheat in California, 561–562
Orange, 492
Orange rusts of Rubus, 61–74; of blackberry, 508
ORTON, C. R., see WALTON, R. C.
ORTON, W. A., and BEATTIE, R. KENT, The biological basis of foreign plant quarantines, 295–306
OSMUN, A. V., and ANDERSON, P. J., An improved formaldehyde tank for the onion drill, 58; see ANDERSON, P. J.
Oversummering, of cereal leaf rust, 338–340
Oxalate, presence in fungous filtrates, 549
Oxygen relations, Urocystis tritici, 133–134; Puccinia coronata, 133

PAINE, SYDNEY G., and CHAUDHURI, H., The blackleg disease of the potato. On the relationship of Bacillus atrosepticus and Bacillus solanisaprus, 359–361
Panax quinquefolium, host of Heterodera radicicola, 442
Papaya, Helminthosporium on, 484; see Carica papaya
Panicum, anceps, clandestinum, dichotomiflorum, and gattingeri, hosts for zonate eye-spot, 59
Para rubber tree, host of Helminthosporium heveae, 483

Paris daisy, host of Bacterium tumefaciens, 87–89

Paris daisy, host of Bacterium tumefaciens, 87–89

PARKER, C. S., Coryneum blight of stone fruits in Washington, 510; Notes on the anthracnose of lettuce, 510

PAXTON, G. E., see MACKIE, W. W.

Pea, intracellular bodies in tissues of, 326–329; root rot and blight of, 293

Peach, aerial galls of, 45; host of Coryneum beijerinckii, 510; Polystictus hirsutus on, Polystictus lacteus on, 185

Pear, fruit rot due to Botrytis cinerea, 509

Pecan scab, Fusicladium effusum, 300

Pectic fermentation, by Bacillus caratovorus, Bacterium angulatum, B. campestre, B. sojae, and B. tabacum, 381–384

Penicillium, susceptibility of to copper, 539

Pennisetum japonicum, host for zonate eye-spot, 59

Pennsylvania, apple blotch infection in 1922 in, 43

Peony, blight due to Botrytis cinerea, and B. paeoniae, 509; diseases, 292; blight by Botrytis paeoniae, 292; blotch by Cladosporium paeoniae, 292; leaf-spot by Septoria paeoniae berolinensis, 292; mosaic, 292; leaf, petal and stem spot of, by Septoria paeoniae berolinensis, 290

Pepper, seed treatment in diseases of, 57; bacterial spot of, 307–315

Periwinkle, leaf blight due to Botrytis sp., 509

Personals, 148–149; 289; 410; 464; 563

Pestalozzia, relation of an undescribed species to a disease of the camphor tree, 47; guepinii, 486; guepinii vaccinii, causing cranberry rot, 510

Pfeffer's nutrient agar, nematode infection in, 439

Phaeolus sistotremoides, 294

Phalaris arundinacea, host for zonate eye-spot, 59

Phaseolus, max, host of Heterodera radicicola, 443; radiatus, host of Heterodera radicicola, 442; vulgaris, anthracnose of, 316–323

Philippines, gum diseases on sugar cane in, 504

Phleum pratense, host for zonate eye-spot, 59; host of Ustilago striaeformis, 38

Phlyctaena linicola, disease of flax, 53

Phoma, 461–462; stage of Mycosphaerella rubina on raspberries, 44; destructiva, pepper disease, 58; lingam, hot water treatment in cabbage seed, 251–253

Phomopsis, sp., causing cranberry rot, 510; sulphur toxic to spp., 60; citri, melanose of citrus fruits, 45; juniperovora, in relation to damping-off of conifers, 387–391; pseudotsuga, on Pseudotsuga, Larix, Tsuga, and Abies, 301

Photography, photographing spore discharge in apothecia, 30–32

Phyllachora trifolii, 118

Phyllosticta, congesta, on plum, 462

Physiology, of perithecial and pycnidial formation in Valsa leucostoma, 257–272

Phytopathological notes, 145–148, 187, 283–289, 410, 461–464, 504–506

Phytopathologists, notice of international conference of, in Holland, June 25 to 30, 1923, 287, 288

Phyllosticta solitaria, infection in 1922 in Pennsylvania, 43–44

Physalis spp., as mosaic carrier, 555; heterophylla, host of tomato mosaic, 372–375; minima, host of Bacterium vesicatorium, 311, 314; pubescens, host for potato mosaic, 41; subglabrata, overwintering of tomato mosaic on, 372–375; virginiana, host of tomato mosaic, 372

Physiology, of plant pathogenic bacteria, 381–384

Phytophthora, evidence indicating that it should be merged with Pythium, 34; allii, 529; arecae, 523, 529; cactorum, 526; capsici, 529; colocasiae, 529; cryptogea on tomato, 520, 529; erythroseptica, 524, 525, 529; infestans on tomatoes, 520, 529; on Ito's variety of potato, 56; melongonae, 529; **mexi-**

cana, n. sp., on tomato, 520–531; technical description of, 529–530; parasitica, 527, 529; phaseoli, 524, 529–530; terrestria on tomato, 520, 527, 529
Picea engelmanni, damping-off of, 385
Pigeon pea, wilt of, 543
Pink root, of onion, 290, 291, 508
Pinus albicaulis, host for white pine blister rust of Europe, 45; balfouriana, doubtfully infected with white pine blister rust in Europe, 45; banksiana, damping-off of, 385; koraiensis, host for white pine blister rust, 45; monticola, 511; host of Fomes pinicola, 511; host for white pine blister rust in Northwest, 46; resinosa, damping-off of, 385; strobiliformis, inoculated with white pine blister rust, 45; strobus, foreign studies of white pine blister rust, 45; host for white pine blister rust in Northwest, 46
Piptoporus suberosus, 294
Pisum sativum, host of Heterodera radicicola, 443
Plant, disease survey of England and Wales, 410; diseases of 1922 in western Quebec, 290; quarantines, foreign, biological basis of, 295–306
Plasmopara viticola, on grapes, 518; susceptibility of to copper, 539
Platanus, 185
Pleosphaerulina, 117; briosiana, 122
Plum, blotch of, 461–462; fruit rot due to Botrytis cinerea, 509; Polystictus hirsutus on, 185; Polystictus lacteus on, 185
Poa pratensis, host of Septoria nodorum, and Septoria tritici, 18; host for zonate eye-spot, 59; host of timothy rust, 38; Septoria leaf blotch of, 11–13
Polyporus, barbaeformis, see Poria papyracea; fissus, 294
Polystictus, abietinus, on Abies grandis, 184; on Abies balsamea, 184; on Thuja plicata, 184; on Tsuga heterophylla, 184; biformis, on Fagus atropunicea, 184; on Quercus rubra, 184; cervinus, see P. biformis; cinnabarinus, on Betula occidentalis, 184–186; conchifer, on Ulmus spp., 184; floridanus, on Chamaecyparis thyoides, 184; on Liquidambar styraciflua, 184; hirsutus, on plum, cherry, prune, peach and apple, 185; on Acer rubrum, 185; on Alnus tenuifolia, 185; on Betula occidentalis, 185; on Castanea dentata, 185; on Quercus coccinea, 185; on Q. prinus, 185; lacteus, on plum, cherry, peach and apple, Acer, Platanus, Quercus and Ailanthus, 185; pargamenus, on Acer saccharum, 185; on Betula lutea, 185; on Fraxinus americana, 185; on Nyssa sylvatica, 185; on Populus tremuloides, 185; on Quercus alba, 185; on Q. coccinea, 185; on Q. marilandica, 185; pinsitus, on Chamaecyparis thyoides, 185; on Juniperus virginiana, 185; versicolor, 185; zonatus, on Populus, sp., 185; on Salix sp., 185
Polystictus and decay of living trees, 184–186
Populus, eugenei, Dothichiza populea on, in Ohio, 245–247; tremuloides, Polystictus pargamenus on, 185
Poria, lindbladii, see Polystictus floridanus; papyracea, on Chamaecyparis thyoides, 187; on Thuja occidentalis, 187; subacida, 294
Porodaedalea pini, 294
Portulaca oleracia, host of Heterodera radicicola, 442
Potato, agar, formula of, 544; mosaic diseases of, 555–557; Alternaria blight of, 100; black leg disease, 359–361; causes of stem-end discolorations, 55; Ito's variety in New York, 55; diseases of, 40; marginal leaf roll, 40; spindling tuber, 40; Dutch varieties resistant to wart disease and internal brown spot, 54; host of Bacterium vesicatorium, 307–315; trypanosome-like bodies in tissues of, 324–325; hot water and hot air treatment, 55, 465–475; mosaic masking at high altitudes, 39; mosaic and leaf roll in the Northwest, 39; presprinkling before seed treatment, 512; results of treatment for Rhizoctonia, 509; results of experiments with leaf roll and mosaic in Idaho, 511; scab

in new land, 241–244; hopper burn of, 237–241; scab, sulphur for the control of, 512; relation of soil moisture to, 231–236; seed treatments, 341–358; thermal death point, 465–475; spraying and dusting of, in 1922, in Michigan, 54
Powdery mildew, of raspberry, 508
Precipitated sulphur, 507
Protozoa, occurrence in plants affected with mosaic, 41
Prune, Polystictus hirsutus on, 185
Prunus, americana, susceptibility to Aphis attack, 438; besseyi, immunity to Aphis, 438; hortulana, 462; persica, host of Heterodera radicicola, 443; triflora, 462
Pseudotsuga, sp., host of Phomopsis pseudotsugae, 301; taxifolia, Fomes pinicola on, 511
Puccinia, antirrhini, reaction to fungicides, 532–542; coronata, oxygen relations, 133; dispersa, on rye, 49; graminis and barberry in Denmark, 404–407; graminis avenae, biologic specialization of, 35; graminis secalis, biologic specialization of, 35; graminis tritici, susceptibility of F_2 hybrids of Kanred and Marquis, 437; simplex, on barley, 49; triticina, strains in United States, 36
Purslane, see Portulaca oleracea
Pyropolyporus robiniae, 294
Pyrus malus, Fusicoccum on, 225–230
Pythium, evidence indicating that Phytophthora should be merged with it, 34; artotrogus, in relation to damping-off of conifers, 387–391; related to blossom-end decay of watermelons, 57; deBaryanum, relation of temperature to the damping-off of garden cress seedlings, 273–282; in relation to damping-off of conifers, 386–391; watermelon decay, 57

Quack grass, Septoria leaf blotch of, 7–8
QUANJER, H. M., see BOTJES, J. CORTWIN
Quercus, alba, Polystictus pargamenus on, 185; coccinea, host of Polyporus spraguei, 288; Polystictus hirsutus on, 185; marylandica, host of Polyporus spraguei, 288; Polystictus pargamenus on, 185; prinus, host of Polyporus spraguei, 288; Polystictus pargamenus on, 185; Polystictus hirsutus on, 185; rubra, host of Polyporus spraguei, 288; Polystictus biformis on, 184; velutina, host of Polyporus spraguei, 288

RAEDER, J. M., Effect of presprinkling with water on the hot formaldehyde and corrosive sublimate methods of potato seed treatment, 512; Preliminary results with the use of sulphur for the control of potato scab in Idaho, 512
RAMSEY, G. B., see LINK, GEO. L. L.
Raspberry, black, blue stem disease of, 293; mosaic of, control of by rogueing, 292; Phoma stage of Mycosphaerella rubina on, 44–45; diseases, 507; yellows, 507, mosaic, 508; eastern blue-stem, 507, 508, leaf curl, 507, 508; rogueing for, 507; powdery mildew, 508; fruit rot due to Botrytis cinerea, 509
RATHBUN, ANNIE E., Damping-off of taproots of conifers, 385–391
Red clover, see Trifolium pratense
REDDICK, DONALD, Ito's potato variety Ekishirazu in New York, 55–56
REDDY, CHAS. S., and GODKIN, JAMES, A bacterial disease of brome grass, 75–86
Report of the fourteenth annual meeting of the American Phytopathological Society, 188–198
REPPERT, R. R., Cartoons in Agricultural teaching, 33
Resistance to nematodes, 436–450
Rheosporangium, aphanidermatus, in relation to damping-off of conifers, 387–391
Rhizoctonia, disease of flax, 53; notes on, 509; potato seed treatment for, 509; potomacensis, in relation to damping-off of conifers, 387–391; solani (Corticium vagum) in relation to damping-off of conifers, 387–391; sp., on lettuce, 508
Rhizopus, relation of humidity to infection of sweet potato by, 56; nigricans, susceptibility to fungicides, 533–542
RHOADS, ARTHUR S., Notes on the failure of grapevines to set fruit and on shelling, 513–519

Ribes, americanum, host of Cronartium ribicola, 285; aureum, host of C. ribicola, 285; cynosbati, host of C. ribicola, 285; spraying of against rust, 534; nigrum, host of C. ribicola, 283, 285; vulgare, host of C. ribicola, 285; spp., attacked by white pine blister rust in Michigan, 285
Rice, Helminthosporium disease, 53
Richard's solution, composition of, 544
RICHARDSON, J. K., Bacterial soft rot of iris, 293
RIKER, A. J., The location of the crown-gall organism in its host tissues, 43; Some morphological responses of the host tissues to the crown-gall organisms, 43
ROBERTS, JOHN W., A method of isolating selected single spores, 558–560; Apple bitter rot cankers in the eastern United States, 461; Plum blotch, 461–462
Robinia pseudo-acacia, slime bodies in, 327–328, 332–333
Rogueing for raspberry diseases, 507
ROLDAN, EMILIANO, F., see WELLES, COLIN G.
Root rot, of canning peas, 293; strawberry, 508; Texas, recent studies, 33
Rose, blossom blight due to Botrytis sp., 509; host of Bacterium tumefaciens, 87–99
Rosette, of raspberry, 507; of wheat intracellular bodies associated with, 41
Rot, due to Botrytis cinerea, on apple, carrot, cranberry, pear, plum, raspberry, strawberry, squash, 509; of lettuce, due to Botrytis, 508; onion, by Botrytis allii, 509; onions, due to Fusarium, 510; soft, of iris, 293; stem-end, of citrus fruits, 45
Rots, cranberry, report on, 509–510
Rubber, nurseries, Helminthosporium disease in, 484; tree, Helminthosporium heveae on, 483–487
Rubus spp., hosts of orange rust, 63–74
Rust, of asparagus, 300; of apple, 303; of cedar, 303; black stem, severity of attacks in Denmark, 1884 to 1921, 404–407; leaf rust of barley and rye, 49–50; leaf rust of wheat in United States, 36; orange, of blackberry, 508; of sunflower, 290; white pine blister, 283; in the Northwest, 46; infection through grafted roots, 46; foreign studies of, 45; in Michigan, 285, 286
Rye, leaf rust of, 49; occurrence of bunt in, 210–215; Septoria leaf blotch of, 1–2; host of Septoria secalis, 1–2; host of Septoria nodorum, 18; host of Septoria tritici, 18

Saccharum, spontaneum, host of Sclerospora Philippinensis, 299
Salix, 185
SANFORD, G. B., The relation of soil moisture to the development of common scab of potato, 231–236
Scab, apple, control in Massachusetts, 44; of apple, 508; citrus, history of, in Japan, 492–495; of pecan, 300; of potato, relation of soil moisture to, 231–236; of potatoes in new land, 241–244; potato, use of sulphur for the control of, 512
SCHMITZ, HENRY, Leaf cast of Larix occidentalis by Hypodermella laricis Tubeuf in North Idaho, 505–506; Studies in wood decay V. Physiological specialization in Fomes pinicola Fr., 511
SCHULTZ, E. S., and FOLSOM, DONALD, Spindling tuber and other degeneration diseases of Irish potatoes, 40
Sclerospora, production of conidia at night, 34; Javanica, 34; graminicola, 34; maydis, 34; on maize, 299; Philippinensis, 34; on maize, 299; on Saccharum spontaneum, 299; on Miscanthus japonicus, 299; sacchari on maize, 299; sacchari, 34; spontanea, 34; on maize, 299
Sclerotinia, disease on sunflower, 496–499; from apple, 462; taxonomy of, on Helianthus annuus, 496; cinerea, sulphur toxic to, 60; libertiana, on dahlias, 285; on lettuce, 508; perplexa, 497; sclerotiorum, 497, 499
Sclerotium compactum Helianthi, 499;

rhizodes, disease of wheat caused by, 463–464
Secale, cereale, occurrence of bunt in, 210–215; Septoria leaf blotch of, 1–2; host of Septoria nodorum, 16; host of Septoria tritici, 16, 18
Seed, cotton, fungi from interior of, 501–503; injury in treatment for control of loose smut of wheat, 38; transmission of lettuce mosaic, 104–106; treatments, for control of barley smuts, 153–160; modified hot water and formaldehyde, 154–156; chlorophol and copper carbonate, 157–160; treatment in pepper diseases, 57–58; with organic mercury compounds, 38
Seed-O-San, 176
Septobasidium, frustulosum, on Fraxinus sp., 187; retiforme, 187
Septoria, disease of rye, barley and certain grasses, 1–23; leaf blotch of rye, 1–2; leaf blotch of barley, 3–7; leaf blotch of quack grass, 7–8; leaf blotch of brome grass, 8–11; of Kentucky blue grass, 11–13; agropyri, on Agropyron repens, 7–8, 18–23; bromi, 21–22; on Bromus altissimus, 9; on Bromus inermis, 8–11; on Bromus secalinus, 9; nodorum, 18–23; on Poa pratensis, 18; on Secale cereale, Triticum aestivum, T. compactum, T. dicoccum, T. duram, 16; T. monococcum, 17; T. polonicum, 17; spelta, 17; T. turgidum, 17; passerinii, 18–23; on Hordeum spp., 3–7; on Hordeum vulgare pallidum, 3; H. deficiens, 15; H. horsfordianum, 15; H. distichon erectum, 15; H. distichon nudum, 15; H. distichon nutans, 15; H. vulgare coerulescens, 15; H. vulgare hexastrichum, 15; H. vulgare himalaya, 15; H. vulgare nigrum, 16; H. vulgare trifurcatum, 16; secalis, 18–23; on rye, 1–2; on Secale cereale, 16; tritici, 18–23; on Poa pratensis, 18; on rye, 18; on Secale cereale, 16; on Triticum aestivum, 16; T. compactum, 16; T. durum, 16; T. monococcum, 17; T. polonicum, 17; T. spelta, 17; T. turgidum, 17
Serrano, F. B., see Lee, H. A.
Sereh disease of sugar cane, in Singapore, 145
Sesamum indicum, host of Helminthosporium sesameum, 53
Sesbania, host of Heterodera radicicola, 442, 443; macrocarpa, host of Heterodera radicicola, 443
Severin, Henry H. P., see Kofoid, Charles A.
Simmonds, P. M., see Fraser, W. P.
Single spore isolation methods, 558–560
Slime bodies in Robinia pseudo-acacia, 327–328, 332–333
Smut, control of, by copper carbonate, 293, 294; control of stinking smut of wheat, 37; of timothy, 38–39; covered, on barley, 551–554, infection studies, 551–554; barley, experiments with hot water, formaldehyde, copper carbonate, and chlorophol for the control of, 153–160; flag, of wheat, 127–139; loose smut, of wheat, treatment for control, 38; of oats, control of, by copper carbonate, 291–292; by formaldehyde, 292; onion, 508; treatment for, 161–168; stinking, 512; timothy smut, germination of spores, 38–39; wheat smut investigations in Kansas, 36
Snowball blossom blight due to Botrytis sp., 509
Soil, moisture, relation of, to formaldehyde injury of onion seedlings, 392–403; relation to scab of potato, 231–236
Soja max, host of Heterodera radicicola, 443
Solanaceae, mosaic diseases of, 555
Solanum, carolinense, as mosaic carrier, 555; dulcamara, host of Bacterium vesicatorium, 311, 314; nigrum, host of Bacterium vesicatorium, 311, 314; rostratum, host of Bacterium vesicatorium, 311, 314
South India, Helminthosporium heveae in, 483
Soybean, see Soja max
Spaulding, Perley, Foreign studies of white pine blister rust, 45; Plant disease survey of England and Wales, 410

Sphaeronema fimbriatum, ascigerous stage of, 56
Sphaerulina, leaf spot of clover, 117–126; trifolii, 117–126; trifolii, leaf-spot of clover, 59
Spinach, wilt of, due to Fusarium **spinaciae**, 205–209
Spindling-tuber, and other diseases of Irish potato, 40
Spore, discharge from apothecia, photographing, 30–32; isolation methods, 558–560; selection, methods of, 558; measurements of Helminthosporium heveae, 485
Sporonema, oxycocci causing cranberry rot, 510; pulvinatum, causing cranberry rot, 510
Spot, internal brown, Dutch varieties of potato resistant to, 54
Spraying and dusting experiments, in orchards, 507; potatoes in 1922 in Mich., 54
Squash, storage rot due to Botrytis cinerea, 509; see Cucurbita moschata
STAKMAN, E. C., LEVINE, M. N., and BAILEY, D. L., Biologic specialization of Puccinia graminis avenae, 35; see LEVINE, M. N.
Staling products of fungi, 33–34; of Botrytis cinerea, 33–34; of Fusarium sp., 33–34
Stemphylium sp., causing cranberry rot, 510
Stem rot, of sunflower, 290
Sterigmatocystis, susceptibility of to copper, 539
STEVENS, F. L., and DOWELL, RUTH I., A Meliola disease of cacao, 247–250
STEVENSON, F. J., see GAINES, E. F.
Stickers, for spray mixtures, 507
Stinking smut, 512; control, 169–183; further notes on the relation of the spore load to, 334–337
Stizolobium deeringianum, host of Heterodera radicicola, 443
STONE, R. E., Root rot and blight of canning peas, 293; see HOWITT, J. E.
Stone fruits, Coryneum blight of, in Washington, 510
Storage rot, of carrot and squash, 509; of cranberries, 509–510
Strawberry, fruit rot due to Botrytis cinerea, 509; leaf-spot by Mycosphaerella fragariae, 508; leaf scorch by Mollisia earliana, 508; root rot of, 508; winter injury of, 508
Streak, of melons, 462
Sugar beets, curly leaf of in Argentine, 458–460
Sugar cane, gum diseases of, in the Philippines and Australia, 504; bacterial diseases, 504; mosaic, transmission by Aphis maidis in Porto Rico, 24–29; red vascular disease, 504; Sereh disease, 145
Sulphur, as a fungicide, 60, 177; for tobacco leaf-spots, 142; colloidal, and precipitated, 507; results on use of, to control potato scab, 512
Sulphuric acid, 501
Sumatra, Helminthosporium heveae in, 483–487
SUMMERBY, R., see DICKSON, B. T.
Summer meeting of the American Phytopathological Society, 507–508
Sunflower, host of Sclerotinia, 496–499; rust, stem rot of, wilt of, 290
Sunscald, cause of, on beans, 376–380
Susceptibility, of Prunus americana to aphis attack, 438
Sweet pea, intracellular bodies in tissues of, 326–329
Sweet potatoes, ascigerous stage of black-rot fungus, 56; mosaic diseases of, 42–43; mottle-necrosis, 145–146; internal break down, 146–147; relation of humidity to infection by Rhizopus, 56
SWEZY, OLIVE, see KOFOID, CHARLES A.

Take-all, of wheat, 508
TANAKA, TYÔZABURÔ, A brief history of citrus scab in Japan, 492–495
Tangelo, 492
Tango, see Chrysanthemum coronarium
Tank, formaldehyde, for onion drill, 161–168
TAPKE, V. F., Modified and simplified hot-water and vapor treatments for the

control of loose smut in wheat, with special reference to seed injury, 38
Tarnish plant bug, 512
TAUBENHAUS, J. J., and KILLOUGH, D. T., Recent studies on control methods of Texas root rot, 33
TAYLOR, J. W., see TISDALE, W. H.
TAYLOR, MINNIE W., White pine blister rust infection through grafted roots, 46; see COLLEY, REGINALD H.
Temperature, in relation to growth of Macrosporium cucumerinum, 199–204; influence on blight of wheat and corn, 50; influence on morphology of Fusarium spores, 51; relation of, to the damping-off of garden-cress seedlings by Pythium deBaryanum and Corticium vagum, 273–282; relations, Urocystis tritici, 130–133; studies in mosaic diseases, 42
Texas root rot, on Ipomoea trichocarpa, 33
Theabromo cacao, Meliola **guianensis,** and Helminthosporium **guianensis,** on 247–250
THOMPSON, NOEL F., Eradicating the common barberry by means of chemicals, 48
Thuja plicata, Polystictus abietinus on, 184; occidentalis, Poria papyracea on, 187
Thurberia thespesioides, host of Bacterium malvacearum, 455–457
Tilletia, tritici, wheat smut investigations in Kansas, 36; laevis, dust treatments for, 37; tritici, cytological studies, 36–37; dusting for, 169–183; investigations of, in Kansas, 36; on rye, 210–215
Time-temperature curves for killing potatoes by heat treatment, 465–475
TIMS, E. C., see JONES, L. R.
Tip burn, lettuce, 508
TISDALE, W. H., An effective method of inoculating barley with covered smut, 551–554; and TAYLOR, J. W., Organic mercury seed disinfectants, 38; TAYLOR, J. W., and GRIFFITHS, MARION A., Experiments with hot water, formaldehyde, copper carbonate, and chlorophol for the control of barley smuts, 153–160
Tobacco, angular leaf spot, 140; controlling wildfire in seed-beds, 59; crown gall, 107–116; dust, 507; leaf-spots, 140–144; mosaic diseases of, 555; Olpidium brassicae on, 452; species of Olpidium parasitic on, 451–454; wildfire, 140
Tomato, blight, 520; buckeye-rot of, 520; collar-rot of, 520; damping-off of, 520; mosaic diseases of, 555; stem girdling of, 520; disease of, caused by Phytophthora mexicana, 520–531; field control of mosaic, 372–375; nematode infection on, 439–440; host of Heterodera radicicola, 442; hybrids, susceptibility to nematodes, 437; lightning injury, 147; mosaic, Nelson's spiral bodies in, 330–331; intracellular bodies in tissues of, 326–329; species of Olpidium parasitic on, 451–454; varieties susceptible to Heterodera radicicola, 444, 446–450; and pepper, bacterial spot of, 307–315
Toxicity studies, 532–542
Transmission, of lettuce mosaic by seeds, 104–106
Tree morning glory, see Ipomoea syringaefolia
TREHERNE, R. C., The pollination of certain vegetable plants by insects (Title only), 290
Trifolium, alexandrium, host of Heterodera radicicola, 443; hybridum, leaf-spot, 59; host of Sphaerulina **trifolii,** 117–126; incarnatum, host of Heterodera radicicola, 442; pratense, host of Heterodera radicicola, 442; host of Sphaerulina **trifolii,** 117–126; leaf spot, 59; intracellular bodies in tissues of, 326–329; repens, host of Heterodera radicicola, 442; host of Sphaerulina **trifolii,** 117–126; leaf-spot, 59
Trigonella foenum graceum, host of Heterodera radicicola, 443
Triticum, aestivum, host of Septoria nodorum, 16; host of Septoria tritici, 16; compactum, host of Septoria nodorum, S. tritici, 16; dicoccum, host of

Septoria nodorum, 16; durum, host of Septoria nodorum, S. tritici, 16; monococcum, host of Septoria nodorum, S. tritici, 17; polonicum, host of Septoria nodorum, S. tritici, 17; spelta, host of Septoria nodorum, S. tritici, 17; turgidum, host of Septoria nodorum, S. tritici, 17; strains of leaf rust, 36

Trypanosome like bodies in Solanaceous plants, 324–325

Tsuga, heterophylla, Fomes pinicola on, 511; Polystictus abietinus on, 184; sp., host of Phomopsis pseudotsugae, 301

Tulip blight, 509

Turnip juice medium, formula of, 545

Tylenchus dipsaci, 508; physiological strains of, 441

Ulmus spp., Polystictus conchifer on, 184

Urocystis, cepulae, on onion, 508; tritici, causing flag smut of wheat, 127–139; relation to temperature, 130–133; morphologic features of germination, 134–137

Uromyces caryophyllinus, reaction to fungicides, 532–542

Ustilago hordei, 153–160; infection studies of, 551–554; levis, 171, 183; nuda, 153–160; striaeformis, germination of spores, 38–39

VALLEAU, W. D., An important period in the life history of two bacterial organisms causing leaf-spots on tobacco, 140–144; Oversummering of leaf rust of cereals in Kentucky, 338–340

Valsa leucostoma, the physiology of perithecial and pycnidial formation, 257–272

Venturia inaequalis, control in Massachusetts, 42; reaction to fungicides, 532–542

Verticillium, albo-atrum, causing blue stem disease of black raspberry, 293; sp., causing black heart of apricot, 216–224

VEVE, R. A., see CHARDON, C. E.

Vicia, agritinense, host of Heterodera radicicola, 443; biennis, host of Heterodera radicicola, 443; dasycarpa, host of Heterodera radicicola, 443; faba, host of Heterodera radicicola, 442, 443; fulgens, host of Heterodera radicicola, 443; hybrida, host of Heterodera radicicola, 443; ludoviciana, host of Heterodera radicicola, 443; ludoviciana, host of Heterdera radicicola, 443; pannonica, host of Heterodera radicicola, 443; peregrina, host of Heterodera radicicola, 443; polyphylla, host of Heterodera radicicola, 443; sativa, host of Heterodera radicicola, 443; stricta, host of Heterodera radicicola, 443; villosa, host of Heterodera radicicola, 443

Vigna unguiculata, host of Heterodera radicicola, 442, 443

Vitis, aestivalis, cinerea, cordifolia, 516; labrusca, and rotundifolia, black rot on, 300; vinifera, host for Guignardia bidwellii, 298; host for Plasmopara viticola, 298; host of Heterodera radicicola, 442

Wales, plant disease survey of England and, 410

WALKER, J. C., The hot water treatment of cabbage seed, 251–253; see JONES, L. R.

WALTON, R. C., and ORTON, C. R., Time of apple blotch infection for 1922 in southern Pennsylvania, 43–44

Wart, Dutch varieties of potatoes resistant to disease, 54; of potato, 295

Watermelon, blossom-end decay, 57

WEBB, R. W., see MCKINNEY, H. H.

WEBER, GEORGE F., Mosaic disease of sweet potatoes, 42; III. Septoria disease of rye, barley and certain grasses, 1–23

WEIMER, G. L., see HARTER, L. L.

WEIR, JAMES R., Polyporus spraguei Berk., cause of heart rot, 288; The genus Polystictus and decay of living trees, 184–186; Poria papyracea (Schw.) Cke., 187

WELLES, COLIN G., A new leaf-spot disease of onion and garlic, 362–365; Leaf-spot of lettuce, 289; and ROLDAN, EMILIANO F.

Another economic host of Bacterium solanacearum, 488–491
WENIGER, WANDA, Pathological morphology of durum wheat grains affected with "black point," 48–49; Studies on the causes of stem-end discolorations of potato tubers in North Dakota, 55
Western hemlock, see Tsuga heterophylla
Western white pine, see Pinus monticola
WESTON, WM. H., JR., The production of conidia at night in species of Sclerospora, 34
Wheat, control of stinking smut of, 37; dusting of, for smut, 169–183; flag smut, 127–139; foot-rot of in California, 561, 562; morphology when affected with "black point," 48–49; sclerotium disease of, 463–464; seed treatment, 169–183; seedling blight of, 50; smut investigations in Kansas, 36; stinking smut of, 334–337; take-all, 508
WHETZEL, H. H., The Alternaria blight of potatoes in Bermuda, 100–103
White clover, see Trifolium repens
White fir, see Abies grandis
White pine, blister rust, 299; present status of, in Canada, 291; in Michigan, 285, 286; wintering over of the blister rust of, 283
Wild tobacco, see Nicotiana rustica
Wildfire, of tobacco, 140–144
Wilt, of Chrysanthemum coronarium, 488; of sunflower, 290, 496
WINELAND, GRACE O., The production in culture of the ascigerous stage of Fusarium moniliforme, 51; see JOHANN, HELEN
Winter injury, strawberry, 508
WOLF, FREDERICK A., Studies on the physiology of some plant pathogenic bacteria; VII. Pectic fermentation in culture media containing pectin, 381–384
Wood decay, studies in, 511
WOOLMAN, H. M., Cytological studies on the infection of wheat seedlings by Tilletia tritici (Bjerk.) Wint., 36

Yellows, celery resistance, 56; raspberry, 507
YOUNG, H. C., Sulphur as a fungicide. I. Colloidal sulfur, 60
YOUNG, V. H., Distribution of Ribes and five-needle pine in the Okanogan Region of Washington and British Columbia, 511

Zucker's Bordeaux mixture, toxicity of, 532–542
ZUNDEL, G. L., see HEALD, F. D.